



On large colonies of rufous horseshoe bats (*Rhinolophus rouxii*), western Sri Lanka

Rhinolophus rouxii and *Hipposideros speoris* are small-sized bats belonging to the Rhinolophidae and Hipposideridae families, respectively (Phillips 1935). *Rhinolophus rouxii* is widely distributed in Sri Lanka, India, Nepal, China and Vietnam (Borissenko & Kruskop 2003, Yapa 2017, Wilson & Mittermeier 2019), whereas *H. speoris* is found in Sri Lanka, India and more recently reported from Myanmar (Yapa 2017, Dar *et al.* 2018, Wilson & Mittermeier 2019). These two species are listed as of Least Concern in both Global and National Red Lists of IUCN (IUCN-MOE 2012). As per the existing distribution records, they are commonly found in the low country: *R. rouxii* up to an elevation of 1,590 m, whereas *H. speoris* is at altitudes ranging up to 1,385 m (Phillips 1935, Molur *et al.* 2002, Yapa & Ratnavira 2013, Kusuminda *et al.* 2013, Kusuminda *et al.* 2018, Wilson & Mittermeier 2019).

Observations on population size at two large permanent roosting sites of *R. rouxii* from Hapitigama and Hapitigamkanda and a new site records for *H. speoris* from Hapitigama, Gampaha District, Western province, Sri Lanka (Fig. 1) are reported and analysed here, and in doing so we highlight the significance of abandoned graphite mines for sustaining bat populations. The locality falls within lowland tropical rainforest vegetation (Gunatilleke & Gunatilleke 1990). The annual mean precipitation in 2019 for the nearest meteorological station of Pasyala was 259 mm (highest=714 mm, lowest=67 mm) (Department of Meteorology, Sri Lanka). A field visit was conducted on 17 January 2021 from 08:56 h to 14:40 h. Both roosting sites were identified as a result of an inquiry carried out in the locality. Bats encountered in the sites were caught using hand nets (depth: 45 cm, diameter: 30 cm, mesh size: 1.5 × 1.5 mm). Identification to species

was done based on Phillips (1935), Corbet & Hill (1992), Bates & Harrison (1997) and Sirinivasulu *et al.* (2010). External measurements of 11 variables (Tables 1, 2) were taken following Sirinivasulu *et al.* (2010) with an Incco-RD10 digital caliper to the nearest 0.1 mm, and weight with a precision digital scale (0.01 × 500 g).




Figure 1. Distribution of *Rhinolophus rouxii* (filled circles) and *Hipposideros speoris* (open circles) in Sri Lanka; the localities where both species have been reported together (sympatric) in squares; the present study site is indicated with a star; Historical locations are based on Phillips 1935, Molur *et al.* 2002, Yapa & Ratnavira 2013, Kusuminda *et al.* 2013, and Kusuminda *et al.* 2018; Map ©: A.A.T. Amarasinghe

Captured bats were kept in cotton bags until the measurements were taken. Morphological characters were documented before the bats were released at the capture site. Ambient temperature and relative humidity were measured using a KTJ- TA 38 digital memory thermo-hygrometer and velocity of the wind by a MS-6252 digital anemometer. Roost surface temperature and body temperature of species found were measured with a Work zone-JHK 6606 Digital infra-red thermometer. Light intensity of the microhabitats was measured using a Digitech-QM 157 light meter. Length measurements of the roosting sites and the ambient surroundings were taken using a BOSCH-GLM-100C digital tape and a meter tape. Geographical coordinates and altitude of roosting sites were determined from readings from a Garmin-Etrex hand-held GPS receiver. Photographs of bats and their microhabitats were taken by a Canon-60D camera with Canon 75–300 mm f/4–5.6 and 18–55mm f/3.5–5.6 lenses.

The size of each colony was studied using an incremental visual encounter method and photographic count method following the standard methodologies of Thomas *et al.* (1979) and Thomas & Laval (1988). Visual counts were made by the authors using red filtered head lamps for minimal disturbance, with coordinated efforts and a reasonable amount of time in order to ascertain the accuracy of the colony size and the abundance of species. Sunrise and sunset times on the date of study were recorded as 6.27 h (GMT +5.30) and 18.12 h (GMT +5.30) respectively. Lunar phase was a waxing crescent (21.3%) moon. Both study sites were abandoned graphite mines located within a geographical distance of 500 m from each other.

Study site A: (7°12'06.2" N, 80°07'36.5" E) a mine situated in an old rubber plantation alongside a 1.8 m wide concrete road, close to human habitation with two prominent entrances (see Entrance 1 and 2 in Fig. 2). The canopy cover near Entrance 1 was 60% (11 m in height), dominated by rubber (*Hevea brasiliensis*), kitul palm (*Caryota urens*), coconut (*Cocos nucifera*) and areca palm (*Areca catechu*) trees.

Entrance 1 (alt. 122 m a.s.l.) was an eight meter long horizontal tunnel which connects to the main chamber of the mine (X in Fig. 2) with a 1.6 × 0.9 m sized opening (Fig. 3A). Even though the entire tunnel area was covered by approximately 0.5 m of water, it was easily accessible by humans. According to the locals, the water body is generated from a natural

spring located underneath Entrance 1 of the mine. During the time of observation (08:56 h) ambient air temperature was 26°C (min 26°C, max 26.7°C) and relative humidity was 82.1% (min 79%, max 84%), velocity of wind was recorded as 0 km/h and the light intensity was 19267.4 lux.




Figure 2. Cross section of the floor plan of the mine in study site A from aerial view (not to scale); dotted area shows the area occupied by bats; X and Y indicate the localities of microclimatic data recorded.

Entrance 2 (alt. 135 m a.s.l.) was a 11.2 m deep vertical hole with a 4.1 × 2.6 m sized opening (Figs. 3B–C), located on the hilly side of the plantation, about 15 m behind Entrance 1. The canopy cover around Entrance 2 was 30–40% and mainly consisted of old rubber trees (maximum height 11 m). At the time of observation (11:39 h) ambient air temperature was 28.9 °C (min 25.9 °C, max 28.9 °C) and relative humidity was 85% (min 79%, max 88%), velocity of wind was recorded as 0.75 km/h and the light intensity was 19288.9 lux.

Social calls of the bats were audible by half way through the tunnel from Entrance 1. Sympatric roosting behaviour of *R. rouxii* (Fig. 6) and *H. speoris* (Fig. 7) was evident and most of the ceilings and vertical walls of the mine, except the tunnel area, were occupied by >2000 bats. About 5% (100 individuals) of the bats were *H. speoris* and the remainder were *R. rouxii*. Body temperature of roosting *R. rouxii* ($n=3$) ranged between 23.9–24.8 °C and *H. speoris* ($n=2$) was 24.1 and 24.6° C. Microclimatic conditions recorded at two places (X and Y in Fig. 2) inside study site A are given in Table 3. The morphometric variables and morphological characteristics of the specimens found in study site A are given in Tables 1, 2, and 5.

Table 1. Morphometric variables for *Rhinolophus rouxii* from study site A (Hapitigama) and site B (Hapitigamkanda), Gampaha District, Sri Lanka with a comparison to Phillips (1935) and Bates & Harrison (1997); measurements in mm; — not available.

Measurements	This study				Philips (1935)	Bates & Harrison (1997)	
	site A		site B				
	Female (n=1)	Female (n=1)	Male (n=1)	Female (n=1)			
head and body length	50.2	53.6	48.76	52.51	54.2–60.0	53.1–60.0	47.0–60.0
ear length	17.6	17.5	14.51	17.22	20.0–22.0	19.1–22.0	14.0–22.0
ear width	10.6	11.2	12.41	12.1	—	—	—
forearm length	49.5	47.3	48.16	49.47	48.3–51.0	47.0–54.0	44.0–49.0
wingspan length	330	320	320	310	298–324	286–317	288–304
3 rd metacarpal	40.5	37.7	36.84	39.98	—	—	34.0–39.0
tail length	22.1	18.9	20.19	23.45	25.7–30.0	27.1–30.0	20.0–30.0
tibia length	23.4	21.7	22.03	23.67	—	—	19.0–23.0
hind foot length	10.1	9.7	8.82	9.91	9.1–11.0	9.1–12.0	8.0–11.0
calcar length	12.8	12.2	13.13	13.77	—	—	—
body weight (g)	13.3	11.2	11.72	12.79	—	—	—

Table 2. Morphometric variables for *Hipposideros speoris* from study site A at Hapitigama, Gampaha District, Sri Lanka with a comparison to Phillips (1935) and Bates & Harrison (1997); measurements in mm; — not available.

Measurements	This study		Philips (1935)	Bates & Harrison (1997)	
	site A				
	Male (n=1)	Male (n=1)			
head and body length	56.7	59.2	55.0–56.0	55.0–60.0	46.0–62.0
ear length	15.1	15.0	16.1–18.0	17.1–20.0	12.0–19.0
ear width	12.6	12.5	—	—	—
forearm length	54.8	52.1	52.0–54.0	52.0–54.0	45.6–54.0
wingspan length	340	340	311–324	314–324	—
3 rd metacarpal	43.0	41.5	—	—	35.0–42.0
tail length	24.3	17.8	24.0–30.0	25.0–30.0	20.0–29.0
tibia length	22.2	22.1	—	—	18.0–23.3
hind foot length	8.8	8.21	8.0–9.0	8.0–9.0	7.0–11.0
calcar length	16.0	13.5	—	—	—
body weight (g)	15.0	11.6	—	—	—

Table 3. Microclimatic conditions recorded at two places from study site A (X and Y, Fig. 2) and two places at site B (P and Q; Fig. 4) inside the mines at Hapitigama and Hapitigamkanda (Gampaha District), respectively

Microclimatic condition	site A		site B	
	X	Y	P	Q
temperature °C	26.2	27.2	27.2	26.9
(min, max) inside mines	(25.9, 27.0)	(25.9, 28.9)	(25.9, 26.0)	(25.9, 36.9)
humidity %	85	86	89	85
(min, max)	(79, 87)	(79, 87)	(86, 89)	(76, 88)
temperature (°C)	24–25	24.1	23.3	25.1
outside mines				
velocity of Wind (Km/h)	0	0	0	0
light Intensity (lux)	1932.1	1808.3	796.5	2368.1



Figure 3. Entrance habitat of mine in site A at Hapitigama, Gampaha District, Sri Lanka; outside views of (A) Entrance 1 and (B) Entrance 2; (C) inside view of Entrance 1; (D) main chamber at X

Study site B: ($7^{\circ}11'54.3''\text{N}$ $80^{\circ}07'43.4''\text{E}$; alt.192 m a.s.l.) a mine located in a secondary forest patch with minimal current anthropogenic disturbance. There were two prominent entrances (Fig. 4) about 10 m apart and several other smaller points of entry. Canopy cover near the entrances was 75% (maximum height about 6 m) dominated by Honduran mahogany (*Swietenia macrophylla*), Ceylon breadfruit (*Artocarpus nobilis*), and Cane (*Calamus rotang*). At the time of observation (14:56 h) ambient air temperature near the entrances was 27.8°C (min 25.9°C , max 36°C) and relative humidity was 82% (min 76%, max 89%). Velocity of wind was recorded as 0 km/h and the light intensity was 12690.6 lux. Entrances were vertical tunnels measuring $2.8 \times 2.8 \times 8.5$ m (Entrance 1; Fig. 5A) and $3.6 \times 3.3 \times 13.2$ m (Entrance 2; Fig. 5C–D) (length \times width \times depth) at the opening. Compared to site A, site B had a complicated structure with several lateral and downward passages that either open to large chambers or end blindly. Microclimatic conditions recorded at two places (P and Q in Fig. 4) inside study site B is given in Table 3




Figure 4. Cross section of the floor plan of the mine in study site B from aerial view (not for scale); dotted area shows the area occupied by bats; P and Q indicate the localities of microclimatic data recorded.

Our calculation estimated around 1,500+ *R. rouxii* individuals. However, the actual population size could be much more because many chambers were difficult to approach for counting. There was no evidence of *H. speoris* roosting in this mine. *R. rouxii* occupied almost all the ceilings and vertical walls except the chamber near Entrance 1 (Fig. 4). The body temperature of *R. rouxii* at site B ranged from $24.1\text{--}24.6^{\circ}\text{C}$ ($n=3$). The morphometric variables and morphological characteristics of the specimens found in study site B are given in Tables 1 and 5.



Figure 5. Entrance habitat of mine in site B at Hapitigamkanda, Gampaha District, Sri Lanka; outside views of (A) Entrance 1 and (B) Entrance 2; (C) inside view of Entrance 2; (D) main chamber at P

Historical records indicate that *R. rouxii* had been found commonly in the low country in Sri Lanka up to an altitude of 1,371 m (Fig. 1) and occasionally at higher elevations of 1,829 m in West Haputale (Philips 1935). There are more recent records of frequent occurrence in Radella, Udupussellawa and Idalgashinna up to an elevation of 1,590 m (Kusuminda *et al.* 2020). They roost in either small or very large colonies in caves, graphite mines, culverts, old houses, wells, temples and hollow trees in primary habitats of moist evergreen, lowland forests (Phillips 1935, Bates & Harrison 1997, Molur *et al.* 2002, Yapa 2017, Kusuminda *et al.* 2013, Kusuminda *et al.* 2018, Wilson & Mittermeier 2019). *Hipposideros speoris* had also been recorded throughout the low country in Sri Lanka up to an elevation of 1,385 m (Fig. 1). They roost, singly, in pairs, and in small or large groups in small crevices, caves, tunnels, temples, graphite mines, and abandoned buildings in primary habitats of dry plains to forested hill sides (Phillips 1935, Bates & Harrison 1997, Molur *et al.* 2002, Kusuminda *et al.* 2013, Yapa 2017, Kusuminda *et al.* 2018, Wilson & Mittermeier 2019). Limited literature is available relating to the distribution and population size of *R. rouxii* and *H. speoris* in the Western Province of Sri Lanka. Kusuminda *et al.* (2013) observed both *R. rouxii* and *H. speoris* in the Pilikuttuwa ancient cave temple, which is approximately 17.5 km from the present study sites. But fewer individuals (*R. rouxii*=1, *H. speoris*=25) were recorded during a 14-month study (February 2011–April 2012), and there was no evidence of permanent roosting sites for *R. rouxii* (Kusuminda *et al.* 2013).

Hence, our observations are new site records for *R. rouxii* and *H. speoris* and also represent the two largest documented colonies of *R. rouxii* in the Western province of Sri Lanka. Present observations on both roosting sites and habitat associations agree with the previous autecological reports. Observations on sympatric behavior between *R. rouxii* and *H. speoris* are also in accordance with previous literature (See Fig 1). According to the existing literature *R. rouxii* has two prominent breeding cycles from April to May and from September to November with gestation periods lasting for 150–160 days (Phillips 1935, Bates & Harrison 1997, Yapa 2017, Wilson & Mittermeier 2019). Occurrence of pregnant females from site B indicates that the breeding season of this colony occurs during April to May.

Table 4. Detailed morphological features of *Rhinolophus rouxii* recorded from site A (Hapitigama) and site B (Hapitigamkanda); — not applicable

Morphological characters	Site A (Hapitigama)		Site B (Hapitigamkanda)	
	Female (Gold colour)	Female (Dark brown colour)	Male (Gold colour)	Pregnant Female (Dark brown colour)
Nose shape	The nose-leaf is prominent, complex and triangular shape; short hairs present.			
Head	Small eyes and lower lips with three small grooves			
Ears	Large and pointed			
Inside the ear	Gold colour short hair present	Dark brown short hair present	Gold colour short hair present	Dark brown short hair present
Antitragus	Present	Present	Present	Present
Hair colour on dorsal area	Dark gold colour hair mixed with pale gold colour hair present	Dark brown colour hair mixed with grey colour hairs present	Dark gold colour hair present	Dark brown colour hair present
Breast (Nipple)	Not well-developed and covered with dark gold colour hair present	Not well-developed and covered with dark brown mixed with grey colour hair present	—	Not Well-developed and Covered with dark brown colour hairs present
Hair colour on abdomen	Dark gold colour hair mixed with pale gold colour hair present	Dark brown colour hair mixed with grey colour hair present	Light gold colour hair present	Light brown colour hair present
Ante-brachial membrane	Present			
Radio metacarpal pouch	Absent			
Wing membrane	Naked			
Forearms	Naked			
1 st -5 th metacarpal	Naked			
1 st phalanx 3 rd -5 th metacarpal	Naked			
2 nd phalanx 3 rd -5 th metacarpal	Naked			
Short hairs on Dorsal and Ventral tibia	Naked			
Hair on inter-femoral membrane	Naked			
Wing attached to	back of the tibia above the ankles			
Few long hairs on hind feet	Gold colour short hair present	Dark brown short hair present	Gold colour short hairs present	Dark brown short hairs present
Calcar	Well-developed			
Tail	Fully webbed (enclosed within interfemoral membrane (IFM))			

Plate 16




Figure 6. A *Rhinolophus rouxii* individual (female) from site A (Hapitigama, Gampaha District, Sri Lanka): (A1) ventral aspect of expanded wings, (A2) facial structure. Photo © A.P. Malsha J. Bandara



Figure 7. A *Hipposideros speoris* individual (male) from site A (Hapitigama, Gampaha District, Sri Lanka): (A1) ventral aspect of expanded wings, (A2) facial structure. Photo © A.P. Malsha J. Bandara

Table 5. Detailed morphological features of *Hipposideros speoris* recorded from site A at Hapitigama, Gampaha District, Western province, Sri Lanka; — not applicable

Morphological characters	Male	Male
	(Dark brownish grey colour)	(Dark blackish grey colour)
Nose shape	The Nose-leaf, is Prominent complex and Square shape	
Head	Small eyes, Three small Supplementary leaf lets present	
Frontal sac	Well-developed	
Ears	Rather small, with rounded tips	
Inside the ear	Grey colour short hairs present	Dark blackish grey short hairs present
Antitragus	Conspicuous Spine like projection about one third from the making the antitragus present	
Hair colour on dorsal area	Dark brownish grey colour hairs present	Dark brownish grey colour hairs present
Hair colour on chest	Brownish grey colour hairs present	Dark blackish grey colour hairs present
Hair colour on abdomen	Light Brownish grey colour hairs present	Light blackish grey colour hairs present
Ante-brachial membrane	Present	
Radio metacarpal pouch	Absent	
Wing membrane	Naked	
Forearms	Naked	
1 st -5 th metacarpal	Naked	
1 st phalanx 3 rd -5 th metacarpal	Naked	
2 nd phalanx 3 rd -5 th metacarpal	Naked	
Short hairs on Dorsal and Ventral tibia	Naked	
Hair on inter-femoral membrane (dorsal & ventral)	Naked	
Wing attached to	the tibia just above the Ankles	
Few hairs on hind feet	Brown colour short hairs present	Brownish grey short hairs present
Calcar	Well-developed	
Tail	Fully webbed (enclosed with Inter-femoral membrane (IFM))	

Even though study site B had a more complicated structure and less anthropogenic pressure, the population size of site A was higher and *H. speoris* was absent at site B. Proximity to water bodies could be a reason for the larger population size and the presence of *H. speoris* at site A. Of all types of roosting places, caves provide the most stable conditions for bats, allowing them to colonize in large numbers (Yapa 2017). Similar to caves, abandoned graphite mines also play a vital role in the life of tropical bats, by maintaining stable microclimatic conditions, which is important for breeding and raising pups (Yapa & Ratnasooriya 2008, Yapa 2017). Humidity of the two study sites was relatively high (85–89%), which is preferred by *R. rouxii* (Wilson & Mittermeier 2019) and the ambient temperature of the roosts was also less than the outside temperature. Yapa & Ratnasooriya (2008) further state that there is a close relationship between the roost type and the colony size with large colonies found in caves. Hence, it is important to conserve

roosting places, especially abandoned graphite mines that help to sustain large populations of bats. Based on the discussions we had with locals, these mine-dwelling bats are frequently hunted as a cheap source of meat using thorny branches. Digana *et al.* (2000) have recorded that a whole population of *Hipposiderous lankadiva* in Bogala mine has disappeared due to human hunting. This is a common threat to bats in Sri Lanka and in the whole Asian region (Mildenstein *et al.* 2016, Yapa 2017, Dar *et al.* 2018). Therefore, it is important to educate the locals in order to minimize the killing of bats (Digana *et al.* 2000). Colony sizes of bats have drastically declined over the past few years due to deforestation, roost destruction and human interaction (Yapa 2017, Mouler 2002). Priority conservation attention should be given to Sri Lankan bats regardless of their global conservation status. Establishing baseline data by carrying out systematic field surveys in unexplored areas will play an important role for conserving bat fauna in the island (Lim 2017).

Acknowledgement

We thank W. Bandara for arranging the field visits, officers of Mirigama police station and several villagers including S. Pushpakumara, G. Bandara, A.R.C. Kumara, and N. Madushan for their invaluable support rendered to us during our field visit. Also, we thank P. Disanayake for the sketches of the study sites and S. Karunarathna for the assistance.

Literature cited

- Bates, P.J.J. and D.L. Harrison (1997). *Bats of the Indian Subcontinent*. Harrison Zoological Museum, Kent: 258pp.
- Borissenko, A.V. and S.V. Kruskop (2003). *Bats of Vietnam and Adjacent Territories: an identification manual*. Zoological Museum of Moscow M. V. Lomonosov State University, Moscow: 211pp.
- Corbet, G.B. and J.E. Hill (1992). *The mammals of the Indomalayan Region: a systematic review*. Natural History Museum Publications, Oxford University Press: 488pp.
- Dar, T.H., M. Kamalakannan, C. Venkataraman, and K. Chandra (2018). New record of *Hipposideros speoris* (Chiroptera: Hipposideridae) from Myanmar hidden in the national zoological collection of the Zoological Survey of India. *Mammalia*, 83 (5): 515–517.
- Digana, P.M.C.B., W.B. Yapa, P.V. Randeniya, and W.D. Ratnasooriya (2000). *Predators of Sri Lankan bats*. *Loris*, 22: 44–48.
- Gunatileke, I.A.U.N. and C.V.S. Gunatileke (1990). Distribution of floristic richness and its conservation in Sri Lanka. *Conservation Biology*, 4 (1): 21–31.
- IUCN-MOE (2012). *The National Red List 2012 of Sri Lanka: Conservation Status of the Fauna and Flora*. Ministry of Environment, Colombo, Sri Lanka: 476pp.
- Kusuminda, T.G.T., W.G.M. Edirisinghe, R.P. Nanayakkara, and N. Vishvanath (2013). Diversity and population status of bats in Pilikuttuwa ancient cave temple in Gampaha District, Sri Lanka. *Asian Journal of Conservation Biology*, 2 (2): 136–143.
- Kusuminda, T.G.T., A. Mannakkara, B.D. Patterson, and W.B. Yapa (2018). Bats in tea plantations in Sri Lanka: Species richness and distribution. *Journal of Bat Research & Conservation*, 11 (1): 96–105.
- Lim, B.K. (2017). *High diversity but low profile: new initiatives in the study of bats in Sri Lanka*. *Loris*, 28 (1&2): 21–24.
- Mildenstein, T., I. Tanshi, and P.A. Racey (2016). Exploitation of bats for bushmeat and medicine. In: Voigt C. and T. Kingston (eds.). *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Springer, Cham: 325–376pp.
- Molur, S., G. Marimuthu, C. Srinivasulu, S. Mistry et al. (eds.) (2002). *Status of South Asian Chiroptera: Conservation Assessment and Management Plan (C.A.M.P.) Workshop Report, 2002*. Zoo Outreach Organisation, CBSG South Asia and WILD, Coimbatore: 320pp.
- Phillips, W.W.A. (1935). *A Manual of the Mammals of Ceylon*. Dulau & Company, London: 373pp.
- Srinivasulu, C., P.A. Racey, and S. Mistry (2010). A key to the bats (Mammalia: Chiroptera) of South Asia. *Journal of Threatened Taxa*, 2 (7): 1001–1076.
- Thomas, D.W. and R.K. Laval (1988). Survey and census methods. In: Kunz, T.H. (ed.). *Ecological and Behavioral Methods for the Study of Bats*. Smithsonian Institution Press, Washington DC: 77–89pp.
- Thomas, D.W., M.B. Fenton, and R.M.R. Barclay (1979). Social behavior of the little brown bat, *Myotis lucifugus* I. mating behavior. *Behavioral Ecology & Sociobiology*, 6: 129–136.
- Wilson, D.E. and R.A. Mittermeier (eds.) (2019). *Handbook of the Mammals of the World*. Vol. 9: Bats. Lynx Editions, Barcelona: 1008pp.
- Yapa, W. (2017). *A Field Guide to the Bats of Sri Lanka*. Dilmah Ceylon Tea Company PLC, Colombo, Sri Lanka: 142pp.
- Yapa, W.B. and W.D. Ratnasooriya (2008). *Ecology and Biology of Sri Lankan Bats*. University of Colombo: 28pp.
- Yapa, A. and G. Ratnavira (2013). *The Mammals of Sri Lanka*. Field Ornithology Group of Sri Lanka, University of Colombo, Colombo, Sri Lanka: 1009pp.
- Yapa, W.B., W.D. Ratnasooriya, H.H. Costa, and R. Ruebsamen (2005). In flight and out flight activity patterns of five species of cave dwelling bats in Sri Lanka. *Journal of Science of the University of Kelaniya*, 2: 41–62.

Submitted: 9 April 2021, Accepted: 10 May 2021
Section Editor: Burton Lim

A.P.M.J. Bandara¹ & G. Edirisinghe²

¹ No 153/3, Parakandeniya, Imbulgoda, Sri Lanka
E-mail: malshabandara83@gmail.com

² Biodiversity Conservation Society, No 150/6, Stanley Thilakarathne Mawatha, Nugegoda 10250, Sri Lanka