

An erratum for the “**Table 3**” (page 32), in the article “Amarasinghe, A.A.T., S. Karunarathna, M. Madawala, and A. de Silva (2021). Two new rupicolous day geckos of the *Cnemaspis alwisi* group (Reptilia: Gekkonidae) from Sri Lanka. *Taprobanica*, 10 (1): 23–38 + pls. 3–8”:

The **Table 3** must be read as follows:

Table 3. Current distribution patterns of the *Cnemaspis* species of the *podihuna* clade in each biogeographic region (revised and modified after Amarasinghe & Karunarathna 2020); “—” not recorded yet.

Biogeographic region (alt. range a.s.l.)	podihuna clade		
	<i>scalpensis</i> group	<i>podihuna</i> group	<i>alwisi</i> group
(1). Wet Zone			
Lowland (0–400 m)	—	<i>C. manoae</i> <i>C. molligodai</i>	<i>C. gunawardanai</i> sp. nov. <i>C. rajakarunai</i>
Rakwana hills (above 400 m)	<i>C. godagedarai</i>	—	—
Central highland submontane (400–900 m)	<i>C. anslemai</i> <i>C. scalpensis</i>	—	<i>C. kohukumburai</i>
Central highland montane (above 900 m)	<i>C. gemunu</i>	—	—
(2). Intermediate Zone			
Lowland (0–400 m)	—	—	<i>C. alwisi</i> <i>C. rammalensis</i>
Uva hills (above 300 m)	—	—	<i>C. hitihamii</i> <i>C. nilgala</i>
Central highland submontane (400–900 m)	<i>C. gemunu</i>	—	—
Knuckles submontane (400–900 m)	<i>C. phillipsi</i>	<i>C. kandambyi</i>	<i>C. punctata</i>
Knuckles montane (above 900 m)	<i>C. phillipsi</i>	—	<i>C. punctata</i>
(3). Dry Zone			
Lowland (0–400 m)	—	<i>C. podihuna</i>	<i>C. gunasekarai</i> sp. nov. <i>C. nilgala</i>

Erratum submitted: 28 May 2021

Erratum published: 22 November 2021

<https://doi.org/10.47605/tapro.v10i2.270>