



FOUR NEW ASTERINACEOUS MEMBERS FROM KERALA, INDIA

Sectional Editor: R. K. Verma

Submitted: 16 May 2011, Accepted: 13 June 2011

V. B. Hosagoudar^{1*}, Jacob Thomas¹ and D.K. Agarwal²

¹ Tropical Botanic Garden & Research Institute, Palode - 695 562, Thiruvananthapuram, Kerala, India

* E-mail: vbhosagoudar@rediffmail.com

² Plant Pathology Division, Indian Agricultural Research Institute, New Delhi 110 012, India

Abstract

This paper gives an account of four new members belonging to the genera *Asterina* and *Cirsosia*, namely, *Asterina aristolochiae*, *Asterina phyllanthi-beddomei*, *Cirsosia hopeae* and *Asterina thunbergiicola* Hansford var. *indica* are described and illustrated.

Key words: black mildews, *Asterina*, *Cirsosia*, Kerala, India

Introduction

Peppara and Neyyar is twin and adjacent wildlife sanctuaries located on the western slope and in the penultimate end of the Western Ghats including the Agastyar peak in Thiruvananthapuram District of Kerala state, include the hotspot area of Agastyamala. These sanctuaries lie between 8° 7' - 8° 53' N and 70° 4' - 77° 17' E, surrounded by Kalakkad and Mundandurai wildlife sanctuaries in the East, Palode and Paruthipalli forest range in the North. Both sanctuaries together have an area of 181 km², with an altitudinal range from 100–1864 m a.s.l., temperature from 16-35 °C, annual rain fall about 2800 mm. Agastiar peak (1864 m), Pongalapara (1500 m) and Chemunji (1000 m) are the continuous hill ranges, located towards the Eastern side of the study area and steeply descend towards western side. Because of the varied

topography, this area is rich in its plant diversity. We have been studying the foliicolous fungi of this region since 1996 and the present paper is the novelties of this study.

Taxonomy

1. *Asterina aristolochiae* sp. nov. (Pl. 2, Fig. 1)
Coloniae amphigenae, plerumque epiphyllae, tenues, ad 3 mm diam., confluentes et patentiae. Hyphae pallide brunneae, undulatae, oppositae vel irregulariter laxe ramosae, laxe reticulatae, cellulae 21-36 x 4-6 µm. Appressoria alternata vel unilateralis, unicellularis, ovata, subglobosa, integra vel sublobata, lata posita, sessilis, 4-12 x 7-12 µm. Thyriothecia laxe dispersa, orbicularis, saepe connata, ad 100 µm diam., stellatim dehiscentes ad centro, crenatae vel fimbriatae ad

margine, hyphae fringiorae flexuosae; asci pauci vel numerosi, globosi, octospori, ad 43 µm diam.; ascosporeae oblongae, conglobatae, brunnae, uniseptatae, constrictus ad septatae, 14-17 x 8-10 µm, parietus echinulatus.

Colonies amphigenous, mostly epiphyllous, thin, up to 3 mm in diameter, confluent and cover almost upper surface of the leaves. Hyphae pale brown, undulate, branching opposite to irregular at wide angles, loosely reticulate, cells 21-36 x 4-6 µm. Appressoria alternate to unilateral, unicellular, ovate, subglobose, entire to sublobate, broad based, sessile, 4-12 x 7-12 µm. Thyriothecia loosely scattered, orbicular, often connate, up to 100 µm in diameter, stellately dehisced at the centre, crenate to fimbriate at the margin, fringed hyphae flexuous; asci few to many, globose, octosporous, up to 43 µm in diameter; ascospores oblong, conglobate, brown, uniseptate, constricted at the septum, 14-17 x 8-10 µm, wall echinulate.

Materials examined: type

On leaves of *Aristolochia tagala* Cham. (Aristolochiaceae); *Cat. no.* HClO 48252; *Loc.* Peppara Wildlife Sanctuary, Thiruvananthapuram, Kerala; *Coll.* Jacob Thomas & Vimalkumar; *Date.* 18-XI-2007. Isotype, *Cat. no.* TBGT 2991.

Asterina heterotropae Nakamura on *Heterotropa hirsutisepala*, from Japan and *Asterina thotteae* Hosagoudar & Hanlin on *Thottea* spp. from India are reported on the family Aristolochiaceae (Katamoto, 1975; Hosagoudar & Hanlin, 1995). However, the present species differs from both in having unicellular appressoria.

2. *Asterina phyllanthi-beddomei* sp. nov. (Pl. 2, Fig. 2)

Coloniae epiphyllae, subdensae, ad 1 mm diam., confluentes. Hyphae flexuosae vel anfractuae, alternata vel irregulariter laxe ramosae, laxe reticulatae, cellulae 28-43 x 3-5 µm. Appressoria alternata vel unilateralis, bi-cellula, subantrorsa vel patentia, recta vel curvula, 9-15 µm longa; cellulae basilares cylindratae vel cuneatae, 2-5 µm longae; cellulae apicales ovatae vel plerumque globosae, 3-5- stellatim lobatae, 7-10 x 9-12 µm. Thyriothecia dispersa, orbicularis, ad 140 µm diam., stellatim dehiscentes ad centro, margine crenatae; asci numerosi, globosi, octospori, ad 38 µm diam.; ascosporeae oblongae, conglobatae, brunnae, uniseptatae, constrictus ad septatae, 16-24 x 7-10 µm, parietus glabrus.

Colonies epiphyllous, subdense, up to 1 mm in diameter, confluent. Hyphae flexuous to crooked, branching alternate to irregular at wide

angles, loosely reticulate, cells 28-43 x 3-5 µm. Appressoria alternate to unilateral, two celled, subantrorse to spreading, straight to curved, 9-15 µm long; stalk cells cylindrical to cuneate, 2-5 µm long; head cells ovate to mostly globose, 3-5-times stellately lobate, 7-10 x 9-12 µm. Thyriothecia scattered, orbicular, up to 140 µm in diameter, stellately dehisced at the centre, margin crenate; asci many, globose, octosporous, up to 38 µm in diam.; ascospores oblong, conglobate, brown, uniseptate, constricted at the septum, 16-24 x 7-10 µm, wall smooth.

Materials examined: type

On leaves of *Phyllanthus beddomei* (Gamble) M. Mohanan (Euporbiaceae); *Cat. no.* HClO 48869; *Loc.* Peppara Wildlife Sanctuary, Thiruvananthapuram, Kerala; *Coll.* Jacob Thomas; *Date.* 27-II-2008. Isotype, *Cat. no.* TBGT 3245.

Two species of the genus *Asterina*, namely, *A. phyllanthicola* Sudama Singh and *A. phyllanthigena* Hosagoudar are known on the host genus *Phyllanthus* (Hosagoudar, 2004; Singh, 1980). However, the present new species differs from both in having typically lobate head cells of the appressoria.

Key to the *Asterina* species known on the genus *Phyllanthus*

1. Appressoria lobate *Asterina phyllanthi-beddomei*
Appressoria entire 2
2. Form mycelial net *A. phyllanthigena*
Not so *A. phyllanthicola*

3. *Asterina thunbergiicola* Hansford indica var. nov. (Pl. 3, Fig. 3)

Differt a var. *thunbergiicola* appressoriis et ascosporis longioribus.

Colonies hypophyllous, thin, crustose, up to 5 mm in diameter, confluent. Hyphae crooked, branching irregular at various angles, loosely reticulate to form a mycelial net, cells 21-34 x 2-5 µm. Appressoria alternate, two celled, straight to curved, plugged around stomata of the host leaf, 12-24 µm long, stalk cells cylindrical, 7-12 µm long; head cells ovate, globose to hamate, subangular, angular, narrowly to deeply lobate, 4-12 x 9-15 µm. Thyriothecia scattered, orbicular, often 1-2 connate, up to 180 µm in diameter, stellately dehisce at the centre and dissolved later, margin crenate; asci few to many, globose, octosporous, up to 30 µm in diam.; ascospores, conglobate, brown, uniseptate, constricted at the septum, 16-22 x 7-10 µm, wall smooth. Pycnothyria similar to thyriothecia, smaller; pycnothyriospores pyriform, brown,

apiculate, broadly rounded at one end and, attenuated and truncate at the other, 16-22 x 9-15 μm , wall smooth.

Materials examined: type

On leaves of *Thunbergia* sp. (Thunbergiaceae); *Cat. no.* HClO 48870; *Loc.* Peppara Wildlife Sanctuary, Thiruvananthapuram, Kerala; *Coll.* Jacob Thomas; *Date.* 28-II-2008. Isotype, *Cat. no.* TBGT 3246.

Asterina thunbergiicola Hansford is known on *Thunbergia chrysops* from Sierra Leone, Uganda (Hansford, 1945). However, the new variety differs from the var. *thunbergiicola* in having longer appressoria and ascospores.

4. *Cirsosia hopeae* sp. nov. (Pl. 3, Fig. 4)

Coloniae epiphyllae, subdensae, ad 2 mm diam. Hyphae rectae, plerumque oppositae acuteque ramosae, laxae reticulatae, cellulae 38-48 x 9-12 μm . Appressoria intercalaris, ovata, saepe leniter lateralis, 9-15 μm diam. Thyriothecia dispersa, ad initio rotundata vel ovata, elongata ad maturitatem cum rima longitudinalis ad centre, 300-470 x 250-300 μm , margine crenatae vel fimbriatae, hyphae fringiorae rectae, arte aggregatae et parallel, non-appressoriatae; asci numerosi, globosi, octospori, 35-44 μm diam.; ascosporae obovatae, conglobatae, uniseptatae, fortiter constrictus ad septatae, cinnameneo brunneae, 22-25 x 11-13 μm , parietus echinulatus. Pycnothyria numerosa, thyriotheciis similis; pycnothyriosporae unicellularis, fortiter brunneae, pyriformes, leniter papillatae, 18-20 x 11-13 μm .

Colonies epiphyllous, subdense, up to 2 mm in diameter. Hyphae straight, branching mostly opposite at acute angles, loosely reticulate, cells 38-48 x 9-12 μm . Appressoria intercalary, ovate, often slightly lateral, 9-15 μm in diam. Thyriothecia scattered, initially round to ovate, elongated at maturity with a longitudinal slit at the centre, 300-470 x 250-300 μm , margin crenate to fimbriate, fringed hyphae straight, closely aggregated and parallel, devoid of intercalary appressoria; asci many, globose, octosporous, 35-44 μm in diam.; ascospores obovate, conglobate, uniseptate, deeply constricted at the septum, cinnamon brown, 22-25 x 11-13 μm , wall echinulate. Pycnothyria many, similar to thyriothecia; pycnothyriospores unicellular, deep brown, pyriform, slightly papillate, 18-20 x 11-13 μm .

Materials examined: type

On the leaves of *Hopea ponga* (Dennst.) Mabb. (Dipterocarpaceae); *Cat. no.* HClO 48846; *Loc.* near

Peppara Wildlife Sanctuary, Thiruvananthapuram, Kerala; *Coll.* Jacob Thomas and Vimalkumar; *Date.* 31-III-2007. Isotype, *Cat. no.* TBGT 3222.

Intercalary appressoria, elliptical to elongated thyriothecia with longitudinal dehiscence are the characteristic of the genus *Cirsosia*. There are five species of the genus *Cirsosia* are known. Of these, *C. areacearum* Hosagoudar & Pillai, *C. globuliferae* (Pat.) Arx. and *C. transversalis* Bat. & Maia are known on Arecaceae (Hosagoudar & Pillai, 1993), while, *C. irregularis* (Sydow) Arx is known on *Vatica obtusifolia* from Philippines (Müller & Arx, 1962). *Cirsosia hopeae* differs from it in having epiphyllous colonies in contrast to the hypophyllous, straight mycelium in contrast to crooked, smaller thyriothecia against 750 x 200-300 μm and smaller ascospores 23-25 x 11-12 against 35-36 x 15-16 μm .

Acknowledgements

We thank Director (TBGRI) for the facilities. We are grateful to Ministry of Environment and Forest, New Delhi for the financial support and to the Forest Department, Govt. of Kerala for the forest permission.

Literature cited

Hansford, C. G., 1945. Contributions towards the fungus flora of Uganda VII. New records and revisions. *Proceedings of Linnean Society London*, 157: 20-41.

Hosagoudar, V. B. and C. M. Pillai, 1993. Two interesting *Cirsosia* species on *Calamus* from India. *Mycological Research*, 95: 127-128.

Hosagoudar, V. B., 2004. A new *Asterina* species from Kerala, India. *Zoos' Print Journal*, 19: 1522.

Hosagoudar, V. B. and R. T. Hanlin, 1995. New species of *Asterina* and *Echidnodes* from India. *New Botanist*, 22: 187-192.

Katumoto, K., 1975. The Hemisphaeriales in Japan. *Bulletin of the faculty of agriculture, Yamaguti University*, 26: 45-122.

Müller, E. and J. A. V. Arx, 1962. Die Gattungen der didymosporen Pyrenomyceten. *Beiträge zur Kryptogamenflora der Schweiz*, 11: 1-922.

Singh, S., 1980. *Asterina phyllanthicola* sp. nov. from India. *Transactions of the British Mycological Society*, 74: 204-205.

PLATE 03




Fig. 03: *Asterina thunbergiicola* Hansf. var. *indica* Hosagoudar, Thomas & Agarwal var. nov.
a. Appressoriate mycelium, **b.** Thyriothecium, **c.** Ascus, **d.** Ascospores




Fig. 04: *Cirrosia hopeae* Hosagoudar, Thomas & Agarwal sp. nov.
a. Appressoriate (intercalary) mycelium, **b.** Thyriothecium, **c.** Ascus, **d.** Ascospores